

Part I

Geometry of chaos

WE START OUT with a recapitulation of the basic notions of dynamics. Our aim is narrow; we keep the exposition focused on prerequisites to the applications to be developed in this text. We assume that the reader is familiar with dynamics on the level of the introductory texts mentioned in remark 1.1, and concentrate here on developing intuition about what a dynamical system can do. It will be a broad stroke description, since describing all possible behaviors of dynamical systems is beyond human ken. While for a novice there is no shortcut through this lengthy detour, a sophisticated traveler might bravely skip this well-trodden territory and embark upon the journey at chapter 18.

The fate has handed you a law of nature. What are you to do with it?

1. Define your *dynamical system* (M, f) : the space M of its possible states, and the law f^t of their evolution in time.
2. Pin it down locally—is there anything about it that is stationary? Try to determine its *equilibria / fixed points* (chapter 2).
3. Cut across it, represent as a return map from a section to a section (chapter 3).
4. Explore the neighborhood by *linearizing* the flow; check the *linear stability* of its equilibria / fixed points, their stability eigen-directions (chapters 4 and 5).
5. Does your system have a *symmetry*? If so, you must use it (chapters 10 to 12). Slice & dice it (chapter 13).
6. Go global: train by *partitioning the state space* of 1-dimensional maps. Label the regions by *symbolic dynamics* (chapter 14).
7. Now venture global distances across the system by continuing local tangent space into *stable / unstable manifolds*. Their intersections *partition the state space* in a dynamically invariant way (chapter 15).
8. Guided by this topological partition, compute a set of *periodic orbits* up to a given topological length (chapter 7 and chapter 16).

Along the way you might want to learn about Lyapunov exponents (chapter 6), classical mechanics (chapter 8), and billiards (chapter 9).

Part II

Chaos rules

QUNADRY: all these cycles, but what to do with them? What you have now is a topologically invariant road map of the state space, with the chaotic region pinned down by a rigid skeleton, a tree of *cycles* (periodic orbits) of increasing lengths and self-similar structure. In chapter 18 we shall turn this topological dynamics into a multiplicative operation on the state space partitions by means of transition matrices of chapter 17, the simplest examples of evolution operators. This will enable us to *count* the distinct orbits, and in the process touch upon all the main themes of this book, going the whole distance from diagnosing chaotic dynamics to computing zeta functions.

1. Partition the state space and describe all allowed ways of getting from ‘here’ to ‘there’ by means of transition graphs (transition matrices). These generate the total-ity of admissible itineraries (chapter 17)
2. Learn to count (chapter 18)
3. Learn how to measure what’s important (chapter 19)
4. Learn how to evolve the measure, compute averages (chapter 20)
5. Learn what a ‘Fourier transform’ is for a nonlinear world (chapter 21),
6. and how the short-time / long-time duality is encoded by spectral determinant ex-pression for its spectrum in terms of periodic orbits (chapter 22)
7. Learn how to use short period cycles to describe chaotic world at times much beyond the Lyapunov time (chapter 23)
8. What is all this hard work good for? Deterministic diffusion and foundations of ‘far for equilibrium’ statistical mechanics, for example (chapter 24)
9. Back to hard work: ponder how symmetries simplify spectral determinants (chap-ter 25 and chapter 26)

Part III

Chaos: what to do about it?

WHAT YOU KNOW NOW is to partition topologically and invariantly the state space, compute a hierarchy of cycles, compute spectral determinants and their eigenvalues. What next?

1. Why cycle? (chapter 27)
2. Why does it work? (chapter 28)
3. When does it not work? (chapter 29)
4. What does it have to do with turbulence? (chapter 30)
5. How fat is the turbulent attractor? (chapter 30)
6. There is one that is experimentally relevant (chapter 35)

Part IV

The rest is noise

QUNADRY: all these cycles, but how many do I need? Any physical system suffers background noise, any numerical prediction suffers computational roundoff noise, and any set of equations models nature up to a given accuracy, since degrees of freedom are always neglected. If the noise is weak, the short-time dynamics is not altered significantly: short periodic orbits of the deterministic flow still partition coarsely the state space.

1. What is “noise”? (chapter 37)
2. Variational principles of classical mechanics, and path integrals of quantum mechanics, reimagined (chapter 39)

Part V

Quantum chaos

YOU HAVE MASTERED part II of this book. You can play a game of pinball, and if you are a skilled neuroscientist, you now know how to poke rat brains. You have learned that information about chaotic dynamics can be obtained by calculating spectra of linear operators such as the evolution operator, and that these spectra can be expressed in terms of periodic orbits by means of cycle expansions.

But what happens if we scatter quantum mechanical waves rather than point-like pinballs? Is there a link between quantum-mechanical spectra and the dynamics of the underlying classical flow? The answer is yes, in a very pleasing way - essentially the same ζ functions and cycle expansions describe the classical chaotic dynamics, the stochastic dynamics, and the semiclassical quantum mechanics (chapter 41).

1. We start with a lightning review of quantum mechanics (chapter 42) and then discuss the first semiclassical (or WKB) approach to quantization (chapter 43).
2. Then the semiclassical evolution operator (chapter 44) leads to the semiclassical trace formulas and ζ functions quantization formulas (chapter 45).
3. Their simplest applications are through trace formulas for scattering (chapter 46) and multi-scattering (chapter 47).
4. Now that we have derived the semiclassical weight associated with every unstable periodic orbit, we are now able to put together all ingredients that make the game of pinball unpredictable, and compute a “chaotic” part of the helium spectrum to shocking accuracy (chapter 49).
5. A semiclassical theory in terms of classical dynamics alone cannot be exact. Waves interfere, diffract (chapter 51), and higher \hbar corrections need to be incorporated into the periodic orbit theory.

This part is a collaborative effort of Predrag Cvitanović, Roberto Artuso, Per Dahlqvist, Ronnie Mainieri, Gregor Tanner, Gábor Vattay, Niall Whelan, and Andreas Wirzba.

Part VI

Web appendices

THE CONCEIT of this treatise is to teach you all of classical, stochastic and quantum chaos (or turbulence) in one go. There is so much to say, but when? So many trees are obscuring the grandeur of the forest. So whenever possible, we have moved details of a particular topic into an appendix, numbered as the corresponding chapter of the main text. Were this monograph printed and bound, this part would be left on the ChaosBook.org website, to be consulted if a deeper dive into a particular tangent is desired.

1. A brief history of chaos (appendix [A1](#))
2. Smooth conjugacies (appendix [A2](#))
3. Linear algebra, Hamiltonian Jacobians (appendix [A4](#))
4. Lyapunov exponents done right; transport of vector fields (appendix [A6](#))
5. Discrete symmetries (appendix [A25](#))
6. Cycles (appendix [A16](#))
7. Counting (appendix [A18](#))
8. Implementing evolution (appendix [A19](#))
9. Diffusion (appendix [A24](#))
10. Convergence of spectral determinants (appendix [A28](#))
11. Thermodynamic formalism (appendix [A36](#))
12. Statistical mechanics (appendix [A37](#))
13. Quantum mechanics II (appendix [A45](#))
14. Infinite dimensional operators (appendix [A46](#))

Appendix A1

A brief history of chaos

Laws of attribution

1. **Arnol'd's Law:** everything that is discovered is named after someone else (including Arnol'd's law)
2. **Berry's Law:** sometimes, the sequence of antecedents seems endless. So, nothing is discovered for the first time.
3. **Whiteheads's Law:** Everything of importance has been said before by someone who did not discover it.

— Sir Michael V. Berry

Writing a history of anything is a reckless undertaking, especially a history of something that has preoccupied at one time or other any serious thinker from ancient Sumer to today's Hong Kong. A mathematician, to take an example, might see it this way: “[History of dynamical systems.](#)” Nevertheless, here comes yet another very imperfect attempt.

A1.1 Chaos is born

I'll maybe discuss more about its history when I learn more about it.

— Maciej Zworski

(R. Mainieri and P. Cvitanović)

TRYING TO PREDICT the motion of the Moon has preoccupied astronomers since antiquity. Accurate understanding of its motion was important for determining the longitude of ships while traversing open seas.

Kepler's Rudolphine tables had been a great improvement over previous tables, and Kepler was justly proud of his achievements. He wrote in the introduction to the announcement of Kepler's third law, *Harmonice Mundi* (Linz, 1619) in a style that would not fly with the contemporary *Physical Review Letters* editors:

What I prophesied two-and-twenty years ago, as soon as I discovered the five solids among the heavenly orbits—what I firmly believed long before I had seen Ptolemy's *Harmonics*—what I had promised my friends in the title of this book, which I named before I was sure of my discovery—what sixteen years ago, I urged as the thing to be sought—that for which I joined Tycho Brahé, for which I settled in Prague, for which I have devoted the best part of my life to astronomical contemplations, at length I have brought to light, and recognized its truth beyond my most sanguine expectations. It is not eighteen months since I got the first glimpse of light, three months since the dawn, very few days since the unveiled sun, most admirable to gaze upon, burst upon me. Nothing holds me; I will indulge my sacred fury; I will triumph over mankind by the honest confession that I have stolen the golden vases of the Egyptians to build up a tabernacle for my God far away from the confines of Egypt. If you forgive me, I rejoice; if you are angry, I can bear it; the die is cast, the book is written, to be read either now or in posterity, I care not which; it may well wait a century for a reader, as God has waited six thousand years for an observer.

Then came Newton. Classical mechanics has not stood still since Newton. The formalism that we use today was developed by Euler and Lagrange. By the end of the 1800's the three problems that would lead to the notion of chaotic dynamics were already known: the three-body problem, the ergodic hypothesis, and nonlinear oscillators.

A1.1.1 Three-body problem

Bernoulli used Newton's work on mechanics to derive the elliptic orbits of Kepler and set an example of how equations of motion could be solved by integrating. But the motion of the Moon is not well approximated by an ellipse with the Earth at a focus; at least the effects of the Sun have to be taken into account if one wants to reproduce the data the classical Greeks already possessed. To do that one has to consider the motion of three bodies: the Moon, the Earth, and the Sun. When the planets are replaced by point particles of arbitrary masses, the problem to be solved is known as the three-body problem. The three-body problem was also a model to another concern in astronomy. In the Newtonian model of the solar system it is possible for one of the planets to go from an elliptic orbit around the Sun to an orbit that escaped its dominion or that plunged right into it. Knowing if any of the planets would do so became the problem of the stability of the solar system. A planet would not meet this terrible end if solar system consisted of two celestial bodies, but whether such fate could befall in the three-body case remained unclear.

After many failed attempts to solve the three-body problem, natural philosophers started to suspect that it was impossible to integrate. The usual technique for

integrating problems was to find the conserved quantities, quantities that do not change with time and allow one to relate the momenta and positions at different times. The first sign on the impossibility of integrating the three-body problem came from a result of Bruns that showed that there were no conserved quantities that were polynomial in the momenta and positions. Bruns' result did not preclude the possibility of more complicated conserved quantities. This problem was settled by Poincaré and Sundman in two very different ways [12, 53].

In an attempt to promote the journal *Acta Mathematica*, Mittag-Leffler got the permission of the King Oscar II of Sweden and Norway to establish a mathematical competition. Several questions were posed (although the king would have preferred only one), and the prize of 2500 kroner would go to the best submission. One of the questions was formulated by Weierstrass:

Given a system of arbitrary mass points that attract each other according to Newton's laws, under the assumption that no two points ever collide, try to find a representation of the coordinates of each point as a series in a variable that is some known function of time and for all of whose values the series converges uniformly.

This problem, whose solution would considerably extend our understanding of the solar system, . . .

Poincaré's submission won the prize. He showed that conserved quantities that were analytic in the momenta and positions could not exist. To show that he introduced methods that were very geometrical in spirit: the importance of state space flow, the role of periodic orbits and their cross sections, the homoclinic points.

The interesting thing about Poincaré's work was that it did not solve the problem posed. He did not find a function that would give the coordinates as a function of time for all times. He did not show that it was impossible either, but rather that it could not be done with the Bernoulli technique of finding a conserved quantity and trying to integrate. Integration would seem unlikely from Poincaré's prize-winning memoir, but it was accomplished by the Finnish-born Swedish mathematician Sundman. Sundman showed that to integrate the three-body problem one had to confront the two-body collisions. He did that by making them go away through a trick known as regularization of the collision manifold. The trick is not to expand the coordinates as a function of time t , but rather as a function of $\sqrt[3]{t}$. To solve the problem for all times he used a conformal map into a strip. This allowed Sundman to obtain a series expansion for the coordinates valid for all times, solving the problem that was proposed by Weierstrass in the King Oscar II's competition.

The Sundman's series are not used today to compute the trajectories of any three-body system. That is more simply accomplished by numerical methods or through series that, although divergent, produce better numerical results. The conformal map and the collision regularization mean that the series are effectively in the variable $1 - e^{-\sqrt[3]{t}}$. Quite rapidly this gets exponentially close to one, the radius of convergence of the series. Many terms, more terms than any one has ever

wanted to compute, are needed to achieve numerical convergence. Though Sundman's work deserves better credit than it gets, it did not live up to Weirstrass's expectations, and the series solution did not "considerably extend our understanding of the solar system.' The work that followed from Poincaré did.

A1.1.2 Ergodic hypothesis

The second problem that played a key role in development of chaotic dynamics was the ergodic hypothesis of Boltzmann. Maxwell and Boltzmann had combined the mechanics of Newton with notions of probability in order to create statistical mechanics, deriving thermodynamics from the equations of mechanics. To evaluate the heat capacity of even a simple system, Boltzmann had to make a great simplifying assumption of ergodicity: that the dynamical system would visit every part of the phase space allowed by conservation laws equally often. This hypothesis was extended to other averages used in statistical mechanics and was called the ergodic hypothesis. It was reformulated by Poincaré to say that a trajectory comes as close as desired to any phase space point.

Proving the ergodic hypothesis turned out to be very difficult. By the end of twentieth century it has only been shown true for a few systems and wrong for quite a few others. Early on, as a mathematical necessity, the proof of the hypothesis was broken down into two parts. First one would show that the mechanical system was ergodic (it would go near any point) and then one would show that it would go near each point equally often and regularly so that the computed averages made mathematical sense. Koopman took the first step in proving the ergodic hypothesis when he realized that it was possible to reformulate it using the recently developed methods of Hilbert spaces [98]. This was an important step that showed that it was possible to take a finite-dimensional nonlinear problem and reformulate it as a infinite-dimensional linear problem. This does not make the problem easier, but it does allow one to use a different set of mathematical tools on the problem. Shortly after Koopman started lecturing on his method, von Neumann proved a version of the ergodic hypothesis, giving it the status of a theorem [118]. He proved that if the mechanical system was ergodic, then the computed averages would make sense. Soon afterwards Birkhoff published a much stronger version of the theorem.

chapter 19

A1.1.3 Nonlinear oscillators

The third problem that was very influential in the development of the theory of chaotic dynamical systems was the work on the nonlinear oscillators. The problem is to construct mechanical models that would aid our understanding of physical systems. Lord Rayleigh came to the problem through his interest in understanding how musical instruments generate sound. In the first approximation one can construct a model of a musical instrument as a linear oscillator. But real instruments do not produce a simple tone forever as the linear oscillator does, so Lord Rayleigh modified this simple model by adding friction and more realistic

models for the spring. By a clever use of negative friction he created two basic models for the musical instruments. These models have more than a pure tone and decay with time when not stroked. In his book *The Theory of Sound* Lord Rayleigh introduced a series of methods that would prove quite general, such as the notion of a limit cycle, a periodic motion a system goes to regardless of the initial conditions.

A1.2 Chaos grows up

(R. Mainieri)

The theorems of von Neumann and Birkhoff on the ergodic hypothesis were published in 1912 and 1913. This line of enquiry developed in two directions. One direction took an abstract approach and considered dynamical systems as transformations of measurable spaces into themselves. Could we classify these transformations in a meaningful way? This led Kolmogorov to the introduction of the concept of entropy for dynamical systems. With entropy as a dynamical invariant it became possible to classify a set of abstract dynamical systems known as the Bernoulli systems. The other line that developed from the ergodic hypothesis was in trying to find mechanical systems that are ergodic. An ergodic system could not have stable orbits, as these would break ergodicity. So in 1898 Hadamard published a paper with a playful title of ‘... billiards ...,’ where he showed that the motion of balls on surfaces of constant negative curvature is everywhere unstable. This dynamical system was to prove very useful and it was taken up by Birkhoff. Morse in 1923 showed that it was possible to enumerate the orbits of a ball on a surface of constant negative curvature. He did this by introducing a symbolic code to each orbit and showed that the number of possible codes grew exponentially with the length of the code. With contributions by Artin, Hedlund, and H. Hopf it was eventually proven that the motion of a ball on a surface of constant negative curvature was ergodic. The importance of this result escaped most physicists, one exception being Krylov, who understood that a physical billiard was a dynamical system on a surface of negative curvature, but with the curvature concentrated along the lines of collision. Sinai, who was the first to show that a physical billiard can be ergodic, knew Krylov’s work well.

The work of Lord Rayleigh also received vigorous development. It prompted many experiments and some theoretical development by van der Pol, Duffing, and Hayashi. They found other systems in which the nonlinear oscillator played a role and classified the possible motions of these systems. This concreteness of experiments, and the possibility of analysis was too much of a temptation for [Mary Lucy Cartwright](#) and J.E. Littlewood [25], who set out to prove that many of the structures conjectured by the experimentalists and theoretical physicists did indeed follow from the equations of motion. Birkhoff had found a ‘remarkable curve’ in a two dimensional map; it appeared to be non-differentiable and it would be nice to see if a smooth flow could generate such a curve. The work of Cartwright and Littlewood led to the work of Levinson, which in turn provided the basis for the horseshoe construction of S. Smale.

In Russia, Lyapunov paralleled the methods of Poincaré and initiated the strong Russian dynamical systems school [110]. Andronov carried on with the study of nonlinear oscillators and in 1937 introduced together with Pontryagin the notion of coarse systems. They were formalizing the understanding garnered from the study of nonlinear oscillators, the understanding that many of the details on how these oscillators work do not affect the overall picture of the state space: there will still be limit cycles if one changes the dissipation or spring force function by a little bit. And changing the system a little bit has the great advantage of eliminating exceptional cases in the mathematical analysis. Coarse systems were the concept that caught Smale's attention and enticed him to study dynamical systems.

A1.3 Chaos with us

(R. Mainieri)

In the fall of 1961 Steven Smale was invited to Kiev where he met Arnol'd, Anosov, Sinai, and Novikov. He lectured there, and spent a lot of time with Anosov. He suggested a series of conjectures, most of which Anosov proved within a year. It was Anosov who showed that there are dynamical systems for which all points (as opposed to a non-wandering set) admit the hyperbolic structure, and it was in honor of this result that Smale named these systems Axiom-A. In Kiev Smale found a receptive audience that had been thinking about these problems. Smale's result catalyzed their thoughts and initiated a chain of developments that persisted into the 1970's.

Smale collected his results and their development in the 1967 review article on dynamical systems, entitled "Differentiable dynamical systems" [146]. There are many great ideas in this paper: the global foliation of invariant sets of the map into disjoint stable and unstable parts; the existence of a horseshoe and enumeration and ordering of all its orbits; the use of zeta functions to study dynamical systems. The emphasis of the paper is on the global properties of the dynamical system, on how to understand the topology of the orbits. Smale's account takes you from a local differential equation (in the form of vector fields) to the global topological description in terms of horseshoes.

chapter 15

The path traversed from ergodicity to entropy is a little more confusing. The general character of entropy was understood by Weiner, who seemed to have spoken to Shannon. In 1948 Shannon published his results on information theory, where he discusses the entropy of the shift transformation. Kolmogorov went far beyond and suggested a definition of the metric entropy of an area preserving transformation in order to classify Bernoulli shifts. The suggestion was taken by his student Sinai and the results published in 1959. In 1960 Rohlin connected these results to measure-theoretical notions of entropy. The next step was published in 1965 by Adler and Palis, and also Adler, Konheim, McAndrew; these papers showed that one could define the notion of topological entropy and use it as an invariant to classify continuous maps. In 1967 Anosov and Sinai applied

the notion of entropy to the study of dynamical systems. It was in the context of studying the entropy associated to a dynamical system that Sinai introduced Markov partitions in 1968.

Markov partitions allow one to relate dynamical systems and statistical mechanics; this has been a very fruitful relationship. It adds measure notions to the topological framework laid down in Smale's paper. Markov partitions divide the state space of the dynamical system into nice little boxes that map into each other. Each box is labeled by a code and the dynamics on the state space maps the codes around, inducing a symbolic dynamics. From the number of boxes needed to cover all the space, Sinai was able to define the notion of entropy of a dynamical system. In 1970 Bowen came up independently with the same ideas, although there was presumably some flow of information back and forth before these papers got published. Bowen also introduced the important concept of shadowing of chaotic orbits. We do not know whether at this point the relations with statistical mechanics were clear to everyone. They became explicit in the work of Ruelle. Ruelle understood that the topology of the orbits could be specified by a symbolic code, and that one could associate an 'energy' to each orbit. The energies could be formally combined in a 'partition function' to generate the invariant measure of the system.

After Smale, Sinai, Bowen, and Ruelle had laid the foundations of the statistical mechanics approach to chaotic systems, research turned to studying particular cases. The simplest case to consider is 1-dimensional maps. The topology of the orbits for parabola-like maps was worked out in 1973 by Metropolis, Stein, and Stein [114]. The more general 1-dimensional case was worked out in 1976 by Milnor and Thurston in a widely circulated preprint, whose extended version eventually got published in 1988 [115].

A lecture of Smale and the results of Metropolis, Stein, and Stein inspired Feigenbaum to study simple maps. This led him to the discovery of the universality in quadratic maps and the application of ideas from field-theory to dynamical systems. Feigenbaum's work was the culmination in the study of 1-dimensional systems; a complete analysis of a nontrivial transition to chaos. Feigenbaum introduced many new ideas into the field: the use of the renormalization group which led him to introduce functional equations in the study of dynamical systems, the scaling function which completed the link between dynamical systems and statistical mechanics, and the presentation functions which describe the dynamics of scaling functions.

The work in more than one dimension progressed very slowly and is still far from completed. The first result in trying to understand the topology of the orbits in two dimensions (the equivalent of Metropolis, Stein, and Stein, or Milnor and Thurston's work) was obtained by Thurston. Around 1975 Thurston was giving lectures "On the geometry and dynamics of diffeomorphisms of surfaces." Thurston's techniques exposed in that lecture have not been applied in physics, but much of the classification that Thurston developed can be obtained from the notion of a 'pruning front' formulated independently by Cvitanović.

Once one develops an understanding of the topology of the orbits of a dynamical system, one needs to be able to compute its properties. Ruelle had already generalized the zeta function introduced by Artin and Mazur [2], so that it could be used to compute the average value of observables. The difficulty with Ruelle's zeta function is that it does not converge very well. Starting out from Smale's observation that a chaotic dynamical system is dense with a set of periodic orbits, Cvitanović used these orbits as a skeleton on which to evaluate the averages of observables, and organized such calculations in terms of rapidly converging cycle expansions. This convergence is attained by using the shorter orbits used as a basis for shadowing the longer orbits.

This account is far from complete, but we hope that it will help get a sense of perspective on the field. It is not a fad and it will not die anytime soon.

A1.4 Periodic orbit theory

Pure mathematics is a branch of applied mathematics.

— Joe Keller, asked to define applied mathematics

(P. Cvitanović)

The history of periodic orbit theory is rich and curious; recent advances are equally inspired by more than a century of developments in three separate subjects: 1. *classical chaotic dynamics*, initiated by Poincaré and put on its modern footing by Smale [146], Ruelle [137], and many others, 2. *quantum theory* initiated by Bohr, with the modern 'chaotic' formulation by Gutzwiller [81, 83], and 3. *analytic number theory* initiated by Riemann and formulated as a spectral problem by Selberg [113, 142]. Following different lines of reasoning and driven by different motivations, the three separate roads all arrive at *trace formulas*, *zeta functions* and *spectral determinants*.

The fact that these fields are all related is far from obvious, and even today the practitioners tend to cite papers only from their sub-speciality. In Gutzwiller's words [83], "The classical periodic orbits are a crucial stepping stone in the understanding of quantum mechanics, in particular when then classical system is chaotic. This situation is very satisfying when one thinks of Poincaré who emphasized the importance of periodic orbits in classical mechanics, but could not have had any idea of what they could mean for quantum mechanics. The set of energy levels and the set of periodic orbits are complementary to each other since they are essentially related through a Fourier transform. Such a relation had been found earlier by the mathematicians in the study of the Laplacian operator on Riemannian surfaces with constant negative curvature. This led to Selberg's trace formula in 1956 which has exactly the same form, but happens to be exact." *A posteriori*, one can say that zeta functions arise in both classical and quantum mechanics because the dynamical evolution can be described by the action of linear evolution (or transfer) operators on infinite-dimensional vector spaces. The spectra of these operators are given by the zeros of appropriate determinants. One way to evalu-

section 22.1

ate determinants is to expand them in terms of traces, $\log \det(\mathcal{L}) = \text{tr}(\log \mathcal{L})$. In this way the spectrum of an evolution operator becomes related to its traces, i.e. periodic orbits. A deeper way of restating this is to observe that the trace formulas perform the same role in all of the above problems; they relate the spectrum of lengths (local dynamics) to the spectrum of eigenvalues (global eigenstates), and for nonlinear geometries they play a role analogous to the one that Fourier transform plays for the circle.

exercise 4.1

Distant history is easily sanitized and mythologized. As we approach the present, our vision is inevitably more myopic; for very different accounts covering the same recent history, see V. Baladi [11] (a mathematician’s perspective), and M. V. Berry [18] (a quantum chaologist’s perspective). We are grateful for any comments from the reader that would help make what follows fair and balanced.

M. Gutzwiller was the first to demonstrate that chaotic dynamics is built upon unstable periodic orbits in his 1960’s work on the quantization of classically chaotic quantum systems, where the ‘Gutzwiller trace formula’ gives the semiclassical quantum spectrum as a sum over classical periodic orbits [78–81]. Equally important was D. Ruelle’s 1970’s work on hyperbolic systems, where ergodic averages associated with natural invariant measures are expressed as weighted sums on the infinite set of unstable periodic orbits embedded in the underlying chaotic set [131, 132]. This idea can be traced back to the following sources: 1. the foundational 1967 review [146], where S. Smale proposed as “a wild idea in this direction” a (technically incorrect, but prescient) zeta function over periodic orbits, 2. the 1965 Artin-Mazur zeta function for counting periodic orbits [2], and 3. the 1956 Selberg number-theoretic zeta functions for Riemann surfaces of constant curvature [142]. That one could *compute* using these infinite sets was not clear at all. Ruelle [137] never attempted explicit computations, and Gutzwiller only attempted to implement summations over anisotropic Kepler periodic orbits by treating them as Ising model configurations [82] (In retrospect, Gutzwiller was lucky; it turns out that the more periodic orbits one includes, the worse convergence one gets [26]).

chapter 45

chapter 22

remark 22.2

chapter 18

For a long time the convergence of such sums bedeviled the practitioners, until the mathematically rigorous spectral determinants for hyperbolic deterministic flows, and the closely related semiclassical exact Gutzwiller Zeta functions were recast in terms of highly convergent *cycle expansions*. Under these circumstances, a relatively few short periodic orbits lead to highly accurate long time averages of quantities measured in chaotic dynamics and of spectra for quantum systems. The idea, in a nutshell, is that long orbits are shadowed by shorter orbits, and the n th term in a cycle expansion is the difference between the shorter cycles estimate of the period n -cycles’ contribution and the exact n -cycles sum. For unstable, hyperbolic flows, this difference falls off exponentially or super-exponentially [138]. Contrary to what some literature says, cycle expansions are no more ‘clever resummations’ than the Plemelj-Smithies cumulant evaluation of a determinant is a ‘resummation’, and their theory is considerably more reassuring than what practitioners of quantum chaos fear: there is no ‘abscissa of absolute convergence’, there is no ‘entropy barrier’, and the exponential proliferation of cycles is not the problem.

chapter A46

Cvitanović derived ‘cycle expansions’ in 1986-87, in an effort to prove that the mode-locking dimension for critical circle maps discovered by Jensen, Bak and Bohr [92] is universal; the same kind of periodic orbits are involved in the Hénon map, but now in renormalization ‘time’. The symbolic dynamics of the Hénon attractor (the pruning front conjecture [43]) is coded by transition graphs, topological entropy is given by roots of their determinants. This observation led to the study of convergence of spectral determinants for both discrete-time (iterated maps) and continuous-time deterministic flows (both ODEs and PDEs). Cycle expansions thus arose not from temporal dynamics, but from studies of scalings in period-doubling and cycle-map renormalizations [5, 28, 44]. This work was done in collaboration with R. Artuso (PhD 1987-1989), G. Gunaratne, and E. Aurell (PhD 1984-1989), and it was written under the watchful eye of parrot Gaspar in Fundação de Faca, Porto Seguro, as two long *Recycling of strange sets* papers [4, 5]: *I. Cycle expansions* and *II. Applications*. The main lesson was that one should never split theory and applications into papers numbered I and II; part II, which covers many interesting results, has barely been glanced at by anyone.

chapter 23

chapter 18

chapter 28

The first published paper on these developments was Auerbach *et al.* [7] *Exploring chaotic motion through periodic orbits* (submitted March 1987). Here only a ‘level sum’ approximation (23.39),

section 27.5

$$1 = \sum_{x_j \in \text{Fix } f^n} t_j e^{\beta A(x_j, n)}, \quad t_j = \frac{e^{-n s_{(n)}}}{\Lambda_j}, \quad (\text{A1.1})$$

to the trace formula is presented as an n th order estimate of the leading Perron-Frobenius eigenvalue $s_{(n)}$, and applied to the Hénon attractor (Eq. (4) of the above paper). (The *exact* weight of an unstable prime periodic orbit p (for level sum (21.6)) had been conjectured by Kadanoff and Tang [94] in 1984.) Even as it was written, the heuristics of this paper was rendered obsolete by the exact cycle expansions, and yet, mysteriously, this might be one of the most cited periodic orbits papers.

The first attempt to make cycle expansions accessible to every person was condensed into [Phys. Rev. Letter](#), *Invariant measurement of strange sets in terms of cycles* (submitted March 1988) [36]. However, the two long papers by Artuso *et al.* [4, 5] are a better read.

Several applications of the new methodology are worth mentioning. One was the accurate calculation of the leading dozen eigenvalues of the period-doubling operator [5, 28, 128]. Another breakthrough was the cycle expansion of deterministic transport coefficients [3, 34, 42], such as diffusion constants without *any* probabilistic assumptions. The classical Boltzmann equation for the evolution of 1-particle density is based on *Stosszahlansatz*, the assumption that velocities of colliding particles are not correlated. In periodic orbit theory all correlations are included in cycle averaging formulas, such as the cycle expansion for a particle diffusing chaotically across a spatially-periodic array.

chapter 24

Physicists tend to obsess about matters weightier than iterating maps, so Cvitanović and Eckhardt showed that cycle expansions reproduce quantum resonances

of Eckhardt's 3-disk scatterer [54] to rather impressive accuracy [39] (submitted February 1989). Gaspard and Rice published a lovely triptych of articles (submitted September 1988) about the same 3-disk system (classical, semiclassical and quantum scattering) [67–69]. In 1992 P. E. Rosenqvist [55, 130], in his PhD thesis, combined the magic of spectral determinants with their symmetry factorizations [41, 82] to take cycle expansions to ridiculous accuracy; for example, periodic orbits up to 10 bounces determine the classical escape rate for a 3-disk pinball to be

$$\gamma = 0.4103384077693464893384613078192\dots$$

Try to extract this from a direct numerical simulation, or a log-log plot of level sums (A1.1)! Prior to cycle expansions, the best accuracy that Gaspard and Rice achieved by applying Markov approximations to the spectral determinant [68] was 1 significant digit, $\gamma \simeq 0.45$.

A 3-disk billiard is exceptionally nice, uniformly hyperbolic repeller. More often than not, good symbolic dynamics for a given flow is either not available, or its grammar is not finite, or the convergence of cycle expansions is affected by non-hyperbolic regions of state space. In those cases truncations such as the *stability cutoff* of Dahlqvist and Russberg [47, 48] and Dettmann and Morriss [52] might be helpful. The idea is to truncate the cycle expansion by including only the shadowing combinations of pseudo-cycles $\{p_1, p_2 \dots, p_k\}$ such that $|\Lambda_{p_1} \dots \Lambda_{p_k}| \leq \Lambda_{\max}$, with the cutoff Λ_{\max} equal to or smaller than the most unstable Λ_p in the data set.

chapter 29

section 23.7

It is pedagogically easier to motivate sums over periodic orbits by starting with discrete time dynamical systems, but most flows of physical interest are continuous in time. The weighted averages of periodic orbits for continuous time flows were introduced by Bowen, who treated them as Poincaré section suspensions weighted by the ‘time ceiling’ function, and were incorporated into dynamical zeta functions by Parry and Pollicott [125] and Ruelle [133]. For people steeped in quantum mechanics it all looked very unfamiliar, so in 1991 Cvitanović and Eckhardt reformulated spectral determinants for continuous time flows along the lines of Gutzwiller's derivation of the semi-classical trace formula [40]. As a consequence, quantum mechanicians [18, 96, 99] tend to cite this paper as the first paper on cycle expansions.

chapter 21

2D billiards are only toys, but quantization of helium is surely not just a game. By implementing cycle expansions in 1991, the group of Dieter Wintgen obtained a surprisingly accurate helium spectrum [58, 155] from a small set of its shortest cycles. This happened 50 years after old quantum theory had failed to do so and 20 years after Gutzwiller first introduced his quantization of chaotic systems [81].

The Copenhagen group gave many conference and seminar talks about cycle expansions. In December 1986, Cvitanović presented results on the periodic-orbit description of the topology of Lozi and Hénon attractors and the periodic-orbit computation of associated dynamical averages, at the meeting on “*Chaos and Related Nonlinear Phenomena: Where do we go from here?*” This meeting was organized by Moshe Shapiro and Itamar Procaccia and held in the kibutz Kiryat

section 15.4

Anavim. A great meeting, and Celso Grebogi was in the audience. After the “Where do we go from here?” meeting, the Maryland group wrote a series of papers on unstable periodic orbits, or ‘UPOs’. In the first paper [75], *Unstable periodic orbits and the dimensions of multifractal chaotic attractor* (submitted September 1987), the focus was on fractal dimensions of chaotic attractors, as was the fashion in the late 1980’s. They prove that the natural measure ρ_0 of a mixing hyperbolic attractor is given by the limit of a sum over the unstable periodic points x_j of long period n , embedded in a chaotic attractor. Each periodic point is weighted by the inverse of the product of its periodic orbit’s expanding Floquet multipliers Λ_j , Eq. (14) in their paper:

remark 5.1

$$\rho_0(\mathcal{M}_S) = \lim_{n \rightarrow \infty} \sum_{x_j \in \text{Fix} f^n} \frac{1}{\Lambda_j}, \quad x_j \in \mathcal{M}_S. \quad (\text{A1.2})$$

This is an approximate level sum formula for natural measure, a special case of (A1.1), with leading Perron-Frobenius eigenvalue $s = 0$ (no escape), and $\beta = 0$ (observable = 1). The first paper does cite Auerbach *et al.* [7], in which the same approximate level sum seems to have been published for the first time. Ever since then, various cyclist teams cite exclusively their own papers and some of the mathematicians of the 1970’s.

So you have now written a paper that uses periodic orbits. What is one to cite? Work by Sinai-Bowen-Ruelle is smarter and more profound than the vast majority of ‘chaos’ publications from the 1980s on. If you are not actually computing anything using periodic orbits and are reluctant to refer to recent contributions, you can safely credit Ruelle [132, 137] for deriving the dynamical (or Ruelle) zeta function, and Gutzwiller for formulating semiclassical quantization as a Zeta function over unstable periodic orbits [81, 83]. There are no cycle expansions in these papers or in Bowen’s work (see, for example, the description in [Scholarpedia.org](https://en.wikipedia.org/wiki/Scholarpedia)). If you have computed something using sums weighted by periodic-orbit weights, cite the first paper that introduced them, as well as a useful up-to-date reference, which in this case is [ChaosBook.org](https://chaosbook.org). Do not faint because this web-book is available on (gasp!) the internet - it’s third millennium, and having a continuously updated, hyperlinked and reliable reference has its virtues.

Depending on the context, one should also cite 1) Zoldi and Greenside [156] for being the second to determine unstable periodic orbits (127 of them) for Kuramoto-Sivashinsky, on a domain larger than what was studied by Christiansen *et al.* [27], 2) López *et al.* [107] for being the first to determine relative periodic orbits in a spatio-temporal PDE (complex Landau-Ginzburg), and 3) Kazantsev [95] for being the first to determine periodic orbits in a weather model, and for his variational method for finding periodic orbits. We love these authors, but not for their ‘escape-time weighting’.

remark 23.1

While derivations of (A1.1) by Kadanoff and Tang 1984 and Auerbach *et al.* 1987 were heuristic, Grebogi, Ott and Yorke 1987 prove (A1.2) by taking the $n \rightarrow \infty$ limit. In actual computations it would be madness to attempt to take such limit, as longer and longer periodic orbits are exponentially more and more unstable, exponentially growing in number, and non-computable; and the natural

measure ρ_0 is everywhere singular, with support on a fractal set, with its $n \rightarrow \infty$ limit even more impossible to compute. And why would one take this limit? The whole point of cycle expansions is that it is smarter to compute averages *without* constructing ρ_0 .

Taking a limit to obtain a proof is good mathematics, but in statistical mechanics a partition function is not a limit of anything; it is the full sum of all states. Likewise, its ergodic theory cousin, the spectral determinant is not a long-time limit; it is the exact sum over all periodic orbits. Cycle expansions were introduced in a non-rigorous manner, on purpose [36]: the exposition was meant not to frighten a novice, innocent of Borel measurable α to Ω sets. This was set right in the elegant PhD thesis of H. H Rugh's in 1992, *The correlation spectrum for hyperbolic analytic maps* [138], which proves that the zeros of spectral determinants are indeed the Ruelle-Pollicott resonances [127, 134, 135]. The proof is well within mathematicians' comfort zone, so they tend to cite Rugh's paper as the paper on 'Fredholm determinants', and, as always, throw in "a sense of Grothendieck" for good measure [11, 65], without citing earlier papers on cycle expansions.

chapter 28

If you intend to determine and use periodic orbits, here is the message: Heuristic 'level sums' are approximations to the exact trace formulas (that are derived here, in ChaosBook, and Gaspard monograph [66] with no more effort than the heuristic approximations), not smart for computations; faster convergence is obtained by utilizing the shadowing that is built into the exact cycle expansions of dynamical zeta functions and spectral determinants. Cycle expansions are *not* heuristic, in classical deterministic dynamics they are *exact* expansions in the unstable periodic orbits [4, 5, 36]; in quantum mechanics and stochastic mechanics they are semi-classically exact. So why would one prefer a limit of a heuristic sum such as (A1.2) to the exact spectral determinant, convergent exact periodic orbits sums, and exact periodic orbits formulas for dynamical averages of observables? It is not even wrong. Perhaps if one is very fond of baker's maps [122], which, being piecewise linear, have no cycle expansion curvature terms, one does not appreciate the shadowing cancelations built into the spectral determinants and their cycle expansions. That might be the reason why linear thinkers stop at the level sum (A1.2).

section 27.5

A1.5 Dynamicist's vision of turbulence

The past is never dead. It's not even past.

— William Faulkner, *Requiem for a Nun*

(P. Cvitanović and Lennaert van Veen)

The key theoretical concepts that form the basis of dynamical theories of turbulence of chapter 30 are rooted in the work of Poincaré, Hopf, Smale, Ruelle, Gutzwiller and Spiegel. In his 1889 analysis of the three-body problem [126], Poincaré introduced the geometric approach to dynamical systems and methods

that lie at the core of the theory developed here: qualitative topology of state space flows, Poincaré sections, key roles played by equilibria, periodic orbits, heteroclinic connections, and their stable/unstable manifolds.

In a seminal 1948 paper [87], Ebehardt Hopf visualized the function space of allowable Navier-Stokes velocity fields as an infinite-dimensional state space, parameterized by viscosity, boundary conditions and external forces, with instantaneous state of a flow represented by a point in this state space. Laminar flows correspond to equilibrium points, globally stable for sufficiently large viscosity. As the viscosity decreases (as the Reynolds number increases), turbulent states set in, represented by chaotic state space trajectories. Hopf's observation that viscosity causes a contraction of state space volumes under the action of dynamics led to his key conjecture: that long-term, typically observed solutions of the Navier-Stokes equations lie on finite-dimensional manifolds embedded in the infinite-dimensional state space of allowed states. Hopf's manifold, known today as the 'inertial manifold,' is well-studied in the mathematics of spatio-temporal PDEs. Its finite dimensionality for non-vanishing 'viscosity' parameter has been rigorously established in certain settings by Foias and collaborators [64]. Hopf presciently noted that "the geometrical picture of the phase flow is, however, not the most important problem of the theory of turbulence. Of greater importance is the determination of the probability distributions associated with the phase flow". Hopf's call for understanding probability distributions associated with the phase flow has indeed proven to be a key challenge, one in which dynamical systems theory has made the greatest progress in the last half century. In particular, the Sinai-Ruelle-Bowen ergodic theory of 'natural' or SRB measures has played a critical role in understanding dissipative systems with chaotic behavior [21, 137, 144, 146].

Hopf noted "[t]he great mathematical difficulties of these important problems are well known and at present the way to a successful attack on them seems hopelessly barred. However, there is no doubt that many characteristic features of the hydrodynamical phase flow occur in a much larger class of similar problems governed by non-linear space-time systems. In order to gain insight into the nature of hydrodynamical phase flows we are, at present, forced to find and to treat simplified examples within that class." Hopf's call for geometric state space analysis of simplified models first came to fulfillment with the influential Lorenz's truncation [108] of the Rayleigh-Bénard convection state space. The Proper Orthogonal Decomposition (POD) models of boundary-layer turbulence brought this type of analysis closer to physical hydrodynamics [6, 86]. Further significant progress has proved possible for systems such as the 1-spatial dimension Kuramoto-Sivashinsky flow [101, 145], which is a paradigmatic model of turbulent dynamics, as well as one of the most extensively studied spatially extended dynamical systems.

example 2.2

Today, as we hope to have convinced the reader, with modern computation and experimental insights, the way to a successful attack on the full Navier-Stokes problem is no longer "hopelessly barred." We address the challenge in a way Hopf could not divine, employing methodology developed only within the past two decades, explained in depth in this book.

chapter 30

Hopf, however, to the best of our knowledge, never suggested that turbulent flow should be analyzed in terms of ‘recurrent flows’, i.e. time-periodic solutions of the defining PDEs. The story so far goes like this: in 1960 Ed Spiegel was [Robert Kraichnan](#)’s research associate. Kraichnan told him, “Flow follows a regular solution for a while, then another one, then switches to another one; that’s turbulence.” It was not too clear, but Kraichnan’s vision of turbulence moved Ed. In 1962 Spiegel and Derek Moore investigated a set of 3rd order convection equations which seemed to follow one periodic solution, then another, and continued going from periodic solution to periodic solution. Ed told Derek, “This is turbulence!” and Derek said “This is wonderful!” He gave a lecture at Caltech in 1964 and came back very angry. They pilloried him there. “Why is this turbulence?” they kept asking and he could not answer, so he expunged the word ‘turbulence’ from their 1966 paper [117] on periodic solutions. In 1970 Spiegel met Kraichnan and told him, “This vision of turbulence of yours has been very useful to me.” Kraichnan said: “That wasn’t my vision, that was Hopf’s vision.” What Hopf *actually* said and where he said it remains deeply obscure to this very day. There are papers that lump him together with Landau, as the ‘Landau-Hopf’s incorrect theory of turbulence,’ a proposal to deploy incommensurate frequencies as building blocks of turbulence. This was Landau’s guess and was the only one that could be implemented at the time.

The first paper to advocate a periodic orbit description of turbulent flows is thus the 1966 Spiegel and Moore paper [117, 147]. Thirty years later, in 1996 Christiansen *et al.* [27] proposed (in what is now the gold standard for exemplary [ChaosBook.org/projects](#)) that the periodic orbit theory be applied to infinite-dimensional flows, such as the Navier-Stokes, using the Kuramoto-Sivashinsky model as a laboratory for exploring the dynamics close to the onset of spatiotemporal chaos. The main conceptual advance in this initial foray was the demonstration that the high-dimensional (16-64 mode Galérkin truncations) dynamics of this dissipative flow can be reduced to an approximately 1-dimensional return map $s \rightarrow f(s)$, by choosing the unstable manifold of the shortest periodic orbit as the intrinsic curvilinear coordinate from which to measure near recurrences. For the first time for any nonlinear PDE, some 1,000 unstable periodic orbits were determined numerically. What was novel about this work? First, dynamics on a strange attractor embedded in a high-dimensional space was essentially reduced to 1-dimensional dynamics. Second, the solutions found provided both a *qualitative description* and highly accurate *quantitative predictions* for the given PDE with the given boundary conditions and system parameter values.

How is it possible that the theory originally developed for low dimensional dynamical systems can work in the ∞ -dimensional PDE state spaces? For dissipative flows the number of unstable, expanding directions is often finite and even low-dimensional; perturbations along the ∞ of contracting directions heal themselves, and play only a minor role in cycle weights - hence the long-time dynamics is effectively finite dimensional. For a more precise statement, see Ginelli *et al.* [71].

The 1996 project went as far as one could with methods and computation resources available, until 2002, when new variational methods were introduced [45,

103, 104]. Considerably more unstable, higher-dimensional regimes have become accessible [38]. Of course, nobody really cares about Kuramoto-Sivashinsky. It is only a model; it was not until the full Navier-Stokes calculations of Eckhardt, Kerswell and collaborators [59, 85, 152] that the fluid dynamics community started to appreciate that the *dynamical* (as opposed to *statistical*) analysis of wall-bounded flows is now feasible [70].

A1.6 Gruppenpest

How many Tylenols should I take with this?... (never took group theory, still need to be convinced that there is any use to this beyond mind-numbing formalizations.)

— Fabian Waleffe, forced to read chapter 10.

If you are not fan of chapter 10 “Flips, slides and turns,” and its elaborations, you are not alone. Or, at least, you were not alone in the 1930s. That is when the articles by two young mathematical physicists, Eugene Wigner and Johann von Neumann [119], and Wigner’s 1931 Gruppentheorie [154] started *Die Gruppenpest* that plagues us to this very day.

According to John Baez [9], the American physicist John Slater, inventor of the ‘Slater determinant,’ is famous for having dismissed groups as unnecessary to physics. He wrote:

“It was at this point that Wigner, Hund, Heitler, and Weyl entered the picture with their ‘Gruppenpest:’ the pest of the group theory [actually, the correct translation is ‘the group plague’] ... The authors of the ‘Gruppenpest’ wrote papers which were incomprehensible to those like me who had not studied group theory... The practical consequences appeared to be negligible, but everyone felt that to be in the mainstream one had to learn about it. I had what I can only describe as a feeling of outrage at the turn which the subject had taken ... it was obvious that a great many other physicists were disgusted as I had been with the group-theoretical approach to the problem. As I heard later, there were remarks made such as ‘Slater has slain the ‘Gruppenpest’’. I believe that no other piece of work I have done was so universally popular.”

A. John Coleman writes in *Groups and Physics - Dogmatic Opinions of a Senior Citizen* [29]: “The mathematical elegance and profundity of Weyl’s book [Theory of Groups and QM] was somewhat traumatic for the English-speaking physics community. In the preface of the second edition in 1930, after a visit to the USA, Weyl wrote, “It has been rumored that the ‘group pest’ is gradually being cut out of quantum physics. This is certainly not true in so far as the rotation and Lorentz groups are concerned; ...” In the autobiography of J. C. Slater, published in 1975, the famous MIT physicist described the “feeling of outrage” he and other physicists felt at the incursion of group theory into physics at the hands of Wigner, Weyl et al. In 1935, when Condon and Shortley published their highly influential treatise on the “Theory of Atomic Spectra”, Slater was widely heralded as having

“slain the Gruppenpest”. Pages 10 and 11 of Condon and Shortley’s treatise are fascinating reading in this context. They devote three paragraphs to the role of group theory in their book. First they say, “We manage to get along without it.” This is followed by a lovely anecdote. In 1928 Dirac gave a seminar, at the end of which Weyl protested that Dirac had said he would make no use of group theory but that in fact most of his arguments were applications of group theory. Dirac replied, “I said that I would obtain the results without previous knowledge of group theory!” Mackey, in the article referred to previously, argues that what Slater and Condon and Shortley did was to rename the generators of the Lie algebra of $SO(3)$ as “angular momenta” and create the feeling that what they were doing was physics and not esoteric mathematics.”

From [AIP Wigner interview](#): AIP: “In that circle of people you were working with in Berlin, was there much interest in group theory at this time?” WIGNER: “No. On the opposite. Schrödinger coined the expression, ‘Gruppenpest’ must be abolished.” “It is interesting, and representative of the relations between mathematics and physics, that Wigner’s paper was originally submitted to a Springer physics journal. It was rejected, and Wigner was seeking a physics journal that might take it when von Neumann told him not to worry, he would get it into the Annals of Mathematics. Wigner was happy to accept his offer [141].”

You would think it was all up from there for group theory. But no. In the early 1970’s, in the [nonexistent city of Bielefeld](#), writes M. du Sautoy [140]; “The Maoist movement decided that group theory was a reactionary subject of the old regime, and [...] demonstrations erupted outside of the maths department with protesters holding placards demanding ‘No more group theory’. [...] During one demonstration, the students scaled the outside of the building and scrawled ‘Group Theory Department’ on the wall.

A1.7 Death of the Old Quantum Theory

In 1913 Otto Stern and Max Theodor Felix von Laue went up for a walk up the Uetliberg. On the top they sat down and talked about physics. In particular they talked about the new atom model of Bohr. There and then they made the ‘Uetli Schwur.’ If that crazy model of Bohr turned out to be right, then they would leave physics. It did and they didn’t.

— A. Pais, *Inward Bound: of Matter and Forces in the Physical World*

One afternoon in May 1991, Dieter Wintgen is sitting in his office at the Niels Bohr Institute beaming with the unparalleled glee of a boy who has just committed a major mischief. The starting words of the manuscript he has just penned are

The failure of the Copenhagen School to obtain a reasonable . . .

Wintgen was 34 years old at the time, a scruffy kind of guy, always wearing sandals and holed out jeans, the German flavor of a 90's left winger and mountain climber. He worked around the clock with his students Gregor Tanner and Klaus Richter to complete the work that Bohr himself would have loved to have seen done back in 1916: a 'planetary' calculation of the helium spectrum.

Never mind that the 'Copenhagen School' refers not to the old quantum theory, but to something else. The old quantum theory was no theory at all; it was a set of rules bringing some order to a set of phenomena which defied logic of classical theory. The electrons were supposed to describe planetary orbits around the nucleus; their wave aspects were yet to be discovered. The foundations seemed obscure, but Bohr's answer for the once-ionized helium to hydrogen ratio was correct to five significant figures and hard to ignore. The old quantum theory marched on, until by 1924 it reached an impasse: the helium spectrum and the Zeeman effect were its death knell.

Since the late 1890's it had been known that the helium spectrum consists of the orthohelium and parahelium lines. In 1915 Bohr suggested that the two kinds of helium lines might be associated with two distinct shapes of orbits (a suggestion that turned out to be wrong). In 1916 he got Kramers to work on the problem, and he wrote to Rutherford, "I have used all my spare time in the last months to make a serious attempt to solve the problem of ordinary helium spectrum ... I think really that at last I have a clue to the problem." To other colleagues he wrote that "the theory was worked out in the fall of 1916" and of having obtained a "partial agreement with the measurements." Nevertheless, the Bohr-Sommerfeld theory, while by and large successful for hydrogen, was a disaster for neutral helium. Heroic efforts of the young generation, including Kramers and Heisenberg, were of no avail.

For a while Heisenberg thought that he had the ionization potential for helium, which he had obtained by a simple perturbative scheme. He wrote enthusiastic letters to Sommerfeld and was drawn into a collaboration with Max Born to compute the spectrum of helium using Born's systematic perturbative scheme. To a first approximation, they reproduced the earlier calculations. The next level of corrections turned out to be larger than the computed effect. The concluding paragraph of Max Born's classic "Vorlesungen über Atommechanik" from 1925 sums it up in a somber tone [20]:

(...) the systematic application of the principles of the quantum theory (...) gives results in agreement with experiment only in those cases where the motion of a single electron is considered; it fails even in the treatment of the motion of the two electrons in the helium atom.

This is not surprising, for the principles used are not really consistent. (...) A complete systematic transformation of the classical mechanics into a discontinuous mechanics is the goal towards which the quantum theory strives.

That year Heisenberg suffered a bout of hay fever, and the old quantum theory was dead. In 1926 he gave the first quantitative explanation of the helium spectrum. He used wave mechanics, electron spin and the Pauli exclusion principle,

none of which belonged to the old quantum theory. As a result, planetary orbits of electrons were cast away for nearly half a century.

Why did Pauli and Heisenberg fail with the helium atom? It was not the fault of the old quantum mechanics, but rather it reflected their lack of understanding of the subtleties of classical mechanics. Today we know what they missed in 1913-24, the role of conjugate points (topological indices) along classical trajectories was not accounted for, and they had no idea of the importance of periodic orbits in nonintegrable systems.

Since then the calculation for helium using the methods of the old quantum mechanics has been fixed. Leopold and Percival [105] added the topological indices in 1980, and in 1991 Wintgen and collaborators [58, 155] understood the role of periodic orbits. Dieter had good reasons to gloat; while the rest of us were preparing to sharpen our pencils and supercomputers in order to approach the dreaded 3-body problem, they just went ahead and did it. What it took—and much else—is described in this book.

One is also free to ponder what quantum theory would look like today if all this was worked out in 1917. In 1994 Predrag Cvitanović gave a talk in Seattle about helium and cycle expansions to—inter alia—Hans Bethe, who loved it so much that after the talk he pulled Predrag aside and they trotted over to Hans' secret place: the best lunch on campus (Business School). Predrag asked: "Would quantum mechanics look different if in 1917 Bohr and Kramers *et al.* figured out how to use the helium classical 3-body dynamics to quantize helium?"

Bethe was very annoyed. He responded with an exasperated look - in Bethe Deutschinglish (if you have ever talked to him, you can do the voice over yourself):

"It would not matter at all!"

Well, perhaps appendix A45 proves him wrong...

Commentary

Remark A1.1. Notion of global foliations. For each paper cited in dynamical systems literature, there are many results that went into its development. As an example, take the notion of global foliations that we attribute to Smale. As far as we can trace the idea, it goes back to René Thom; local foliations were already used by Hadamard. Smale attended a seminar of Thom in 1958 or 1959. In that seminar Thom was explaining his notion of transversality. One of Thom's disciples introduced Smale to Brazilian mathematician Peixoto. Peixoto (who had learned the results of the Andronov-Pontryagin school from Lefschetz) was the closest Smale had ever come until then to the Andronov-Pontryagin school. It was from Peixoto that Smale learned about structural stability, a notion that got him enthusiastic about dynamical systems, as it blended well with his topological background. It was from discussions with Peixoto that Smale got the problems in dynamical systems that lead him to his 1960 paper on Morse inequalities. The next year Smale published his result on the hyperbolic structure of the non-wandering set. Smale was not the first to consider a hyperbolic point, Poincaré had already done that; but Smale was the first to introduce a global hyperbolic structure. By 1960 Smale was already lecturing on the horseshoe as a structurally stable dynamical system with an infinity of periodic points and promoting his global viewpoint. (R. Mainieri)

Remark A1.2. A brief history of period doubling universality. Mitchell J. Feigenbaum discovered universality in one-dimensional iterative maps in August 1975. Following Feigenbaum's functional formulation of the problem, in March 1976 Cvitanović derived, in collaboration with Feigenbaum, the equation $g(x) = \alpha g(g(x/\alpha))$ for the period doubling fixed point function (not a big step, it is the limit of Feigenbaum functional recursion sequence), which has since played a key role in the theory of transitions to turbulence. The first published report on Feigenbaum's discovery is dated August 1976 (Los Alamos Theoretical Division Annual Report 1975-1976, pp. 98-102, [read it here](#)). By that time the work had become widely known through many seminars Feigenbaum gave in US and Europe. His first paper, submitted to *Advances in Mathematics* in Nov 1976 was rejected. The second paper was submitted to *SIAM Journal of Applied Mathematics* in April 1977 and rejected in October 1977. Finally, J. Lebowitz published both papers [60, 61] without further referee pain (M. J. Feigenbaum, *J. Stat. Phys.* 19, 25 (1978) and 21, 6 (1979)).

A very informative 1976 review by May [112] describes what was known before Feigenbaum's contribution. The geometric parameter convergence was first noted in 1958 by Myrberg [13, 139], and independently of Feigenbaum, by Grossmann and Thomae [76] in 1977 (without noting the universality of δ). The theory of period-doubling universal equations and scaling functions is developed in Kenway's notes of Feigenbaum 1984 Edinburgh lectures [63] (trifle hard to track down). The elegant unstable manifold formulation of universality given in ChaosBook.org is due to Vul, Khanin, Sinai and Gol'dberg [72, 150, 151] in 1982. The most thorough exposition available is the Collet and Eckmann [30] monograph. For a more recent introduction into renormalization theory that starts out with period doubling before moving on to Quantum Field Theory, see Gurau, Rivasseau and Sfondrini [77].

By 1978 Couillet and Tresser [32, 33] have proposed similar equations. Daido [49] has introduced a different set of universal equations. Derrida, Gervois and Pomeau [51] have extracted a great many metric universalities from the asymptotic regime. Grassberger [73] has computed the Hausdorff dimension of the asymptotic attractor. Lorenz [109] and Daido [50] have found a universal ratio relating bifurcations and reverse bifurcations.

If $f(x)$ is not quadratic around the maximum, the universal numbers will be different - see Vilela Mendés [149] and Hu and Mao [89] for their values. According to Kuramoto and Koga [100] such mappings can arise in chemical turbulence. Nonlinear oscillator; quadratic potential with damping and harmonic driving force exhibit cascades of period-doubling bifurcations [102, 116]. Refs. [22–24] compute solutions of the period-doubling fixed point equation using methods of Schöder and Abel, yielding what are so far the most accurate δ and α . See also Weisstein [153].

Since then we have generalized the universal equations to period n -tuplings; constructed universal scaling functions for all winding numbers in circle maps, and established universality of the Hausdorff dimension of the critical staircase. A nice discussion of circle maps and their physical applications is given in refs. [10, 91, 92]. The universality theory for golden mean scalings is developed in refs. [62, 120, 129, 143]. The scaling functions for circle maps are discussed in ref. [46].

The theory would have remained a curiosity, were it not for the beautiful experiment by Libchaber and Maurer [111], and many others that followed. Crucial insights came from Collet and Eckmann [30] and Collet, Eckmann and Koch [31] who explained how the dynamics of dissipative system (such as a viscous fluid) can become 1-dimensional. The experimental and theoretical developments up to 1990's are summarized in reprint collections by P. Cvitanović [37] and B.-L. Hao [84]. We also recommend Hu [88], Crutchfield, Farmer and Huberman [35], Eckmann [56] and Ott [121]. The period-doubling route to turbulence that is by no means the only way to get there; see Eckmann [56] discussion of other routes to chaos.

By 1979 mathematicians also understood that the numerical methods used by Feigenbaum and Cvitanović to solve the universal equations were in fact convergent. Mathematicians did what they do; they attached various names to the equations, and they changed letters around to make the equations unintelligible to physicists. The re-lettering did not stick, but the renamings did. (This remark is based on a Nordita lecture 1982 in Danish prepared together with Mogens Høgh Jensen, and Cvitanović [37]. Ulla Selmer prepared the drawings, Oblivia Kaypro stood for the initial execution.)

Remark A1.3. Levels of ergodicity. In the mid 1970's A. Katok and Ya.B. Pesin tried to use geometry to establish positive Lyapunov exponents. A. Katok and J.-M. Strelcyn carried out the program and developed a theory of general dynamical systems with singularities. They studied uniformly hyperbolic systems (as strong as Anosov's), but with sets of singularities. Under iterations a dense set of points hits the singularities. Even more important are the points that never hit the singularity set. In order to establish some control over how they approach the set, one looks at trajectories that approach the set by some given ϵ^n , or faster.

Ya.G. Sinai, L. Bunimovich and N.I. Chernov studied the geometry of billiards in a very detailed way. A. Katok and Ya.B. Pesin's idea was much more robust: look at the discontinuity set, take an ϵ neighborhood around it. Given that the Lebesgue measure is ϵ^α and the stability grows not faster than $(\text{distance})^n$. A. Katok and J.-M. Strelcyn proved that the Lyapunov exponent is non-zero.

In mid 1980's Ya.B. Pesin studied the dissipative case. Now the problem has no invariant Lebesgue measure. Assuming uniform hyperbolicity, with singularities, and tying together Lebesgue measure and discontinuities, and given that the stability grows not faster than $(\text{distance})^n$, Ya.B. Pesin proved that the Lyapunov exponent is non-zero, and that SRB measure exists. He also proved that the Lorenz, Lozi and Byelikh attractors satisfy these conditions.

In the systems that are uniformly hyperbolic, all trouble is in differentials. For the Hénon attractor, already the differentials are nonhyperbolic. The points do not separate uniformly, but the analogue of the singularity set can be obtained by excising the regions that do not separate. Hence there are 3 levels of ergodic systems:

1. Anosov flow
2. Anosov flow + singularity set: For the Hamiltonian systems the general case is studied by A. Katok and J.-M. Strelcyn, and the billiards case by Ya.G. Sinai and L. Bunimovich. The dissipative case is studied by Ya.B. Pesin.
3. Hénon case: The first proof was given by M. Benedicks and L. Carleson [14–16]. A more readable proof is given in M. Benedicks and L.-S. Young [17].

Remark A1.4. Is the geometry of nature fractal? By 1983 some physicists were starting to learn that there is a thing called “chaos” [106], a thing stressful, nasty and hard to understand (see this book), so they tried to bypass this whole bit of unpleasantness by getting instead an easy, diagnostic number out of it [93]. They were told that Hausdorff dimension is the way to go. Dimension of the canonical $1/3$'s Cantor set can be explained to a school child, so they tried it out for size on many low-dimensional chaotic attractors, and some crazy high-dimensional ones as well. Our all-time favorite (beyond the ‘Untitled 5’ of figure 1.5) was the claim that the dimension of climate is 3.1 (remember, the policy of ChaosBook is not to pump up citation numbers for silly or plainly wrong papers): In *Deterministic chaos: the science and the fiction*, David Ruelle [136] comments: “[...] one should not believe dimension estimates that are not well below $2 \log_{10} N$. [Authors of ...] claim to find a dimension 3.1 for a ‘climatic attractor’ with $N = 500$ data points. [...] The ‘dimensions’ of the order 6 that are obtained are very close to the upper bound $2 \log_{10} N$ permitted by the Grassberger-Procaccia algorithm [74] (N is the length of the time series used, of the order of 103). The ‘proof’ that one has low dimensional dynamics is therefore inconclusive, and the suspicion is that the time evolutions under discussion do not correspond to low-dimensional dynamics. [...] Readers of *The Ultimate Hitchhiker’s Guide to the Galaxy*, that masterpiece of British literature by D. Adams [1], know that a huge supercomputer has answered ‘the great problem of life, the universe, and everything’. The answer obtained after many years of computation is 42. Unfortunately, one does not know to what precise question this is the answer, and what to make of it. It think that what happened is this. The supercomputer took a very long time series describing all it knew about ‘life, the universe, and everything’ and proceeded to compute the correlation dimension of the corresponding dynamics, using the Grassberger-Procaccia algorithm. This time series had a length N somewhat larger than 10^{21} . And you can imagine what happened. After many years of computation the answer came: dimension is approximately $2 \log_{10} \approx 42$.” In 1998 Avnir, Biham, Lidar, and Malcai [8] explored how much support for the fractal self-similarity hypothesis was there, actually. They found that “the majority of the data that was interpreted in terms of fractality in the surveyed Physical Review journals does not seem to be linked (at least in an obvious way) to existing models and, in fact, does not have theoretical backing. Most of the data represent results from nonequilibrium processes. The common situation is this: An experimentalist performs a resolution analysis and finds a limited-range power law with a value of D smaller than the embedding dimension. Without necessarily resorting to special underlying mechanistic arguments, the experimentalist then often chooses to label the object for which she or he finds this power law a ‘fractal’. This is the fractal geometry of nature.” Their plot says it all: the number

of decades (factors of 10) spanned by experimentally derived scaling exponents peaks at 10 (i.e., one decade).

Remark A1.5. Einstein did it? The first hint that chaos is afoot in quantum mechanics was given in a note by A. Einstein [57]. The total discussion is a one sentence remark. Einstein being Einstein, this one sentence has been deemed sufficient to give him the credit for being the pioneer of quantum chaos [83, 148]. We asked about the paper two people from that era, Sir Rudolf Peierls and Abraham Pais; neither had any recollection of the 1917 article. However, Theo Geisel has unearthed a reference that shows that in early 20s Born did have a study group meeting in his house that studied Poincaré’s *Mécanique Céleste* [126]. In 1954 Fritz Reiche, who had previously followed Einstein as professor of physics in Breslau (now Wrocław, Poland), pointed out to J.B. Keller that Keller’s geometrical semiclassical quantization was anticipated by the long forgotten paper by A. Einstein [57]. In this way an important paper written by the physicist who at the time was the president of German Physical Society, and the most famous scientist of his time, came to be referred to for the first time by Keller [97], 41 years later. But before Ian Percival included the topological phase, and Wintgen and students recycled the Helium atom, knowing *Mécanique Céleste* was not enough to complete Bohr’s original program.

Remark A1.6. Berry-Keating conjecture. A very appealing proposal in the context of semiclassical quantization is due to M. Berry and J. Keating [19]. The idea is to improve cycle expansions by imposing unitarity as a functional equation ansatz. The cycle expansions that they use are the same as the original ones described above [4], but the philosophy is quite different; the claim is that the optimal estimate for low eigenvalues of classically chaotic quantum systems is obtained by taking the real part of the cycle expansion of the semiclassical zeta function, cut off at the appropriate cycle length. M. Sieber, G. Tanner and D. Wintgen, and P. Dahlqvist find that their numerical results support this claim; F. Christiansen and P. Cvitanović do not find any evidence in their numerical results. The usual Riemann-Siegel formulas exploit the self-duality of the Riemann and other zeta functions, but there is no evidence of such symmetry for generic Hamiltonian flows. Also from the point of hyperbolic dynamics discussed above, proposal in its current form belongs to the category of crude cycle expansions; the cycles are cut off by a single external criterion, such as the maximal cycle time, with no regard for the topology and the curvature corrections. While the functional equation conjecture is not in its final form yet, it is very intriguing and fruitful research inspiration.

The real life challenge are generic dynamical flows, which fit neither of extreme idealized settings, Smale horseshoe on one end, and the Riemann zeta function on the other.

Remark A1.7. Sources. The tale of appendix A1.7, aside from a few personal recollections, is in large part lifted from Abraham Pais’ accounts of the demise of the old quantum theory [123, 124], as well as Jammer’s account [90]. In August 1994 Dieter Wintgen died in a climbing accident in the Swiss Alps. Remark A1.3 is based on Ya.B. Pesin’s comments.

References

- [1] D. Adams, *The Ultimate Hitchhiker’s Guide to the Galaxy* (Random House, New York, 1979).

- [2] M. Artin and B. Mazur, “On periodic points”, *Ann. Math.* **81**, 82–99 (1965).
- [3] R. Artuso, “Diffusive dynamics and periodic orbits of dynamic systems”, *Phys. Lett. A* **160**, 528–530 (1991).
- [4] R. Artuso, E. Aurell, and P. Cvitanović, “Recycling of strange sets: I. Cycle expansions”, *Nonlinearity* **3**, 325–359 (1990).
- [5] R. Artuso, E. Aurell, and P. Cvitanović, “Recycling of strange sets: II. Applications”, *Nonlinearity* **3**, 361–386 (1990).
- [6] N. Aubry, P. Holmes, J. L. Lumley, and E. Stone, “The dynamics of coherent structures in the wall region of turbulent boundary layer”, *J. Fluid Mech.* **192**, 115–173 (1988).
- [7] D. Auerbach, P. Cvitanović, J.-P. Eckmann, G. Gunaratne, and I. Procaccia, “Exploring chaotic motion through periodic orbits”, *Phys. Rev. Lett.* **58**, 2387–2389 (1987).
- [8] D. Avnir, O. Biham, D. Lidar, and O. Malcai, “Is the geometry of nature fractal?”, *Science* **279**, 39–40 (1998).
- [9] J. Baez, *This Week’s Finds in Mathematical Physics Week 236*, 2006.
- [10] P. Bak, T. Bohr, and M. H. Jensen, “Mode-locking and the transition to chaos in dissipative systems”, *Physica Scripta* **T9**, 50–58 (1985).
- [11] V. Baladi, *Positive Transfer Operators and Decay of Correlations* (World Scientific, Singapore, 2000).
- [12] J. Barrow-Green, *Poincaré and the Three Body Problem* (Amer. Math. Soc., Providence R.I., 1997).
- [13] M. W. Beims and J. A. C. Gallas, “Accumulation points in nonlinear parameter lattices”, *Physica A* **238**, 225–244 (1997).
- [14] M. Benedicks and L. Carleson, “On iterations of $1 - ax^2$ on $(-1, 1)$ ”, *Ann. Math.* **122**, 1–25 (1985).
- [15] M. Benedicks and L. Carleson, On the Hénon attractor, in IXth Int. Congr. on Mathematical Physics, edited by B. Simon, A. Truman, and M. Davies (1989), pp. 498–500.
- [16] M. Benedicks and L. Carleson, “The dynamics of the Hénon map”, *Ann. Math.* **133**, 73 (1991).
- [17] M. Benedicks and L.-S. Young, “Absolutely continuous invariant measures and random perturbations for certain one-dimensional map”, *Ergod. Theor. Dynam. Syst.* **12**, 13–37 (1992).
- [18] M. V. Berry, “Martin Gutzwiller and his periodic orbits”, *Commun. Swiss Phys. Soc.* **37**, 34–38 (2012).
- [19] M. V. Berry and J. P. Keating, “A rule for quantizing chaos”, *J. Phys. A* **23**, 4839–4849 (1990).
- [20] M. Born, *Vorlesungen über Atommechanik*, English Translation: *The Mechanics of the Atom* (Ungar Publishing, New York, 1927).
- [21] R. Bowen, *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms* (Springer, New York, 1975).

- [22] K. Briggs, “A precise calculation of the Feigenbaum constants”, *Mathematics of Computation* **57**, 435–439 (1991).
- [23] K. M. Briggs, T. W. Dixon, and G. Szekeres, “Analytic solutions of the Cvitanović-Feigenbaum and Feigenbaum-Kadanoff-Shenker equations”, *Int. J. Bifur. Chaos* **8**, 347–357 (1998).
- [24] K. M. Briggs, G. R. W. Quispel, and C. J. Thompson, “Feigenvalues for Mandelsets”, *J. Phys. A* **24**, 3363–3368 (1991).
- [25] M. L. Cartwright and J. E. Littlewood, “On non-linear differential equations of the second order”, *J. London Math. Soc.* **20**, 180–189 (1945).
- [26] F. Christiansen and P. Cvitanović, “Periodic orbit quantization of the anisotropic Kepler problem”, *Chaos* **2**, 61–69 (1992).
- [27] F. Christiansen, P. Cvitanović, and V. Putkaradze, “Hopf’s last hope: Spatiotemporal chaos in terms of unstable recurrent patterns”, *Nonlinearity* **10**, 55–70 (1997).
- [28] F. Christiansen, P. Cvitanović, and H. H. Rugh, “The spectrum of the period-doubling operator in terms of cycles”, *J. Phys. A* **23**, L713S–L717S (1990).
- [29] A. J. Coleman, “Groups and physics – Dogmatic opinions of a senior citizen”, *Notices Amer. Math. Soc.* **44**, 8–17 (1997).
- [30] P. Collet and J.-P. Eckman, *Iterated Maps on the Interval as Dynamical Systems* (Birkhäuser, Boston, 2009).
- [31] P. Collet, J.-P. Eckmann, and H. Koch, “Period doubling bifurcations for families of maps on R_n ”, *J. Stat. Phys.* **25**, 1–14 (1981).
- [32] P. Coullet and C. Tresser, “Itérations d’endomorphismes et groupe de renormalisation”, *J. Phys. Colloques C5* **39**, 25–28 (1978).
- [33] P. Coullet and C. Tresser, “Iterations of endomorphisms and renormalization group”, *C. R. Acad. Sc. Paris A* **287**, 577–581 (1978).
- [34] G. Cristadoro, “Fractal diffusion coefficient from dynamical zeta functions”, *J. Phys. A* **39**, L151 (2006).
- [35] J. P. Crutchfield, J. D. Farmer, and B. A. Huberman, “Fluctuations and simple chaotic dynamics”, *Phys. Rep.* **92**, 45–82 (1982).
- [36] P. Cvitanović, “Invariant measurement of strange sets in terms of cycles”, *Phys. Rev. Lett.* **61**, 2729–2732 (1988).
- [37] P. Cvitanović, *Universality in Chaos*, 2nd ed. (Adam Hilger, Bristol, 1989).
- [38] P. Cvitanović, R. L. Davidchack, and E. Siminos, “On the state space geometry of the Kuramoto-Sivashinsky flow in a periodic domain”, *SIAM J. Appl. Dyn. Syst.* **9**, 1–33 (2010).
- [39] P. Cvitanović and B. Eckhardt, “Periodic orbit quantization of chaotic systems”, *Phys. Rev. Lett.* **63**, 823–826 (1989).
- [40] P. Cvitanović and B. Eckhardt, “Periodic orbit expansions for classical smooth flows”, *J. Phys. A* **24**, L237 (1991).
- [41] P. Cvitanović and B. Eckhardt, “Symmetry decomposition of chaotic dynamics”, *Nonlinearity* **6**, 277–311 (1993).

- [42] P. Cvitanović, J.-P. Eckmann, and P. Gaspard, “Transport properties of the Lorentz gas in terms of periodic orbits”, *Chaos Solit. Fract.* **6**, 113–120 (1995).
- [43] P. Cvitanović, G. H. Gunaratne, and I. Procaccia, “Topological and metric properties of Hénon-type strange attractors”, *Phys. Rev. A* **38**, 1503–1520 (1988).
- [44] P. Cvitanović, G. H. Gunaratne, and M. J. Vinson, “On the mode-locking universality for critical circle maps”, *Nonlinearity* **3**, 873–885 (1990).
- [45] P. Cvitanović and Y. Lan, Turbulent fields and their recurrences, in *Correlations and Fluctuations in QCD : Proceedings of 10. International Workshop on Multiparticle Production*, edited by N. Antoniou (2003), pp. 313–325.
- [46] P. Cvitanović, B. Shraiman, and B. Söderberg, “Scaling laws for mode lockings in circle maps”, *Phys. Scr.* **32**, 263–270 (1985).
- [47] P. Dahlgvist, “Determination of resonance spectra for bound chaotic systems”, *J. Phys. A* **27**, 763–785 (1994).
- [48] P. Dahlgvist and G. Russberg, “Periodic orbit quantization of bound chaotic systems”, *J. Phys. A* **24**, 4763–4778 (1991).
- [49] H. Daido, “Universal relation of a band-splitting sequence to a preceding period-doubling one”, *Phys. Lett. A* **86**, 259–262 (1981).
- [50] H. Daido, “Period-doubling bifurcations and associated universal properties including parameter dependence”, *Progr. Theor. Phys.* **67**, 1698–1723 (1982).
- [51] B. Derrida, A. Grevois, and Y. Pomeau, “Universal metric properties of bifurcations of endomorphisms”, *J. Phys. A* **12**, 269–296 (1979).
- [52] C. P. Dettmann and G. P. Morriss, “Stability ordering; strong field Lorentz gas”, *Phys. Rev. Lett.* **78**, 4201–4204 (1997).
- [53] F. Diacu and P. Holmes, *Celestial Encounters: The Origins of Chaos and Stability* (Princeton Univ. Press, Princeton, NJ, 1996).
- [54] B. Eckhardt, “Fractal properties of scattering singularities”, *J. Phys. A* **20**, 5971–5979 (1987).
- [55] B. Eckhardt, G. Russberg, P. Cvitanović, P. E. Rosenqvist, and P. Scherer, “Pinball scattering”, in *Quantum chaos: between order and disorder*, edited by G. Casati and B. Chirikov (Cambridge Univ. Press, Cambridge, 1995), p. 483.
- [56] J.-P. Eckmann, “Roads to turbulence in dissipative dynamical systems”, *Rev. Mod. Phys.* **53**, 643 (1981).
- [57] A. Einstein, “On the quantum theorem of Sommerfeld and Epstein”, in *The Collected Papers of Albert Einstein: The Berlin Years: Writings (1914–1917)*, Vol. 6, E. Schucking, English translation of “Zum Quantensatz von Sommerfeld und Epstein,” *Verh. Deutsch. Phys. Ges.* **19**, 82 (Princeton Univ. Press, Princeton NJ, 1917), p. 443.
- [58] G. S. Ezra, K. Richter, G. Tanner, and D. Wintgen, “Semiclassical cycle expansion for the helium atom”, *J. Phys. B* **24**, L413–L420 (1991).

- [59] H. Faisst and B. Eckhardt, “Traveling waves in pipe flow”, *Phys. Rev. Lett.* **91**, 224502 (2003).
- [60] M. J. Feigenbaum, “Quantitative universality for a class of nonlinear transformations”, *J. Stat. Phys.* **19**, 25–52 (1978).
- [61] M. J. Feigenbaum, “The universal metric properties of nonlinear transformations”, *J. Stat. Phys.* **21**, Reprinted in ref. [37], 669–706 (1979).
- [62] M. J. Feigenbaum, L. P. Kadanoff, and S. J. Shenker, “Quasiperiodicity in dissipative systems: A renormalization group analysis”, *Physica D* **5**, 370–386 (1982).
- [63] M. J. Feigenbaum and R. D. Kenway, The onset of chaos, in *Statistical and Particle Physics: Common Problems and Techniques*. Proc. 26th Scottish Universities Summer School in Physics, edited by K. C. Bowler and A. J. McKane (1984), pp. 1–100.
- [64] C. Foias, B. Nicolaenko, G. R. Sell, and R. Témam, “Inertial manifolds for the Kuramoto-Sivashinsky equation and an estimate of their lowest dimension”, *J. Math. Pure Appl.* **67**, 197–226 (1988).
- [65] D. Fried, “Meromorphic zeta functions for analytic flows”, *Commun. Math. Phys.* **174**, 161–190 (1995).
- [66] P. Gaspard, *Chaos, Scattering and Statistical Mechanics* (Cambridge Univ. Press, Cambridge, 1997).
- [67] P. Gaspard and S. A. Rice, “Exact quantization of the scattering from a classically chaotic repeller”, *J. Chem. Phys.* **90**, 2255–2262 (1989).
- [68] P. Gaspard and S. A. Rice, “Scattering from a classically chaotic repeller”, *J. Chem. Phys.* **90**, 2225–2241 (1989).
- [69] P. Gaspard and S. A. Rice, “Semiclassical quantization of the scattering from a classically chaotic repeller”, *J. Chem. Phys.* **90**, 2242–2254 (1989).
- [70] J. F. Gibson and P. Cvitanović, *Movies of plane Couette*, tech. rep. (Georgia Inst. of Technology, 2011).
- [71] F. Ginelli, P. Poggi, A. Turchi, H. Chaté, R. Livi, and A. Politi, “Characterizing dynamics with covariant Lyapunov vectors”, *Phys. Rev. Lett.* **99**, 130601 (2007).
- [72] A. I. Gol’berg, Y. G. Sinai, and K. M. Khanin, “Universal properties for sequences of bifurcations of period 3”, *Russ. Math. Surv.* **38**, 187–188 (1983).
- [73] P. Grassberger, “On the Hausdorff dimension of fractal attractors”, *J. Stat. Phys.* **26**, 173–179 (1981).
- [74] P. Grassberger and I. Procaccia, “Characterization of strange attractors”, *Phys. Rev. Lett.* **50**, 346–349 (1983).
- [75] C. Grebogi, E. Ott, and J. A. Yorke, “Unstable periodic orbits and the dimensions of multifractal chaotic attractors”, *Phys. Rev. A* **37**, 1711–1724 (1988).
- [76] S. Grossmann and S. Thomae, “Invariant distributions and stationary correlation functions of one-dimensional discrete processes”, *Z. Naturf. A* **32**, 1353–1363 (1977).

- [77] R. Gurau, V. Rivasseau, and A. Sfondrini, [Renormalization: An advanced overview](#), 2014.
- [78] M. C. Gutzwiller, “Phase-integral approximation in momentum space and the bound states of an atom”, [J. Math. Phys.](#) **8**, 1979–2000 (1967).
- [79] M. C. Gutzwiller, “Phase-integral approximation in momentum space and the bound states of an atom. II”, [J. Math. Phys.](#) **10**, 1004–1020 (1969).
- [80] M. C. Gutzwiller, “Energy spectrum according to classical mechanics”, [J. Math. Phys.](#) **11**, 1791–1806 (1970).
- [81] M. C. Gutzwiller, “Periodic orbits and classical quantization conditions”, [J. Math. Phys.](#) **12**, 343–358 (1971).
- [82] M. C. Gutzwiller, “The quantization of a classically ergodic system”, [Physica D](#) **5**, 183–207 (1982).
- [83] M. C. Gutzwiller, [Chaos in Classical and Quantum Mechanics](#) (Springer, New York, 1990).
- [84] B.-L. Hao, [Chaos II](#) (World Scientific, Singapore, 1990).
- [85] B. Hof, C. W. H. van Doorne, J. Westerweel, F. T. M. Nieuwstadt, H. Faisst, B. Eckhardt, H. Wedin, R. R. Kerswell, and F. Waleffe, “Experimental observation of nonlinear traveling waves in turbulent pipe flow”, [Science](#) **305**, 1594–1598 (2004).
- [86] P. Holmes, J. L. Lumley, and G. Berkooz, [Turbulence, Coherent Structures, Dynamical Systems and Symmetry](#) (Cambridge Univ. Press, Cambridge, 1996).
- [87] E. Hopf, “A mathematical example displaying features of turbulence”, [Commun. Pure Appl. Math.](#) **1**, 303–322 (1948).
- [88] B. Hu, “Introduction to real-space renormalization-group methods in critical and chaotic phenomena”, [Phys. Rep.](#) **91**, 233–295 (1982).
- [89] B. Hu and J. M. Mao, “Period doubling: Universality and critical-point order”, [Phys. Rev. A](#) **25**, 3259–3261 (1982).
- [90] M. Jammer, [The Conceptual Development of Quantum Mechanics](#) (McGraw-Hill, New York, 1966).
- [91] M. H. Jensen, P. Bak, and T. Bohr, “Complete devil’s staircase fractal dimension and universality of mode-locking structure in the circle map”, [Phys. Rev. Lett.](#) **50**, 1637–1639 (1983).
- [92] M. H. Jensen, P. Bak, and T. Bohr, “Transition to chaos by interaction of resonances in dissipative systems. I. Circle maps”, [Phys. Rev. A](#) **30**, 1960–1969 (1984).
- [93] L. P. Kadanoff, “Fractals: where’s the physics?”, [Phys. Today](#) **39**, 6 (1986).
- [94] L. Kadanoff and C. Tang, “Escape rate from strange repellers”, [Proc. Natl. Acad. Sci. USA](#) **81**, 1276–1279 (1984).
- [95] E. Kazantsev, “Unstable periodic orbits and attractor of the barotropic ocean model”, [Nonlin. Proc. Geophys.](#) **5**, 193 (1998).

- [96] J. P. Keating, “Resummation and the turning points of zeta functions”, in *Classical, Semiclassical and Quantum Dynamics in Atoms*, edited by B. Eckhardt and H. Friedrich (Springer, Berlin, 1997), pp. 83–93.
- [97] J. B. Keller, “Corrected Bohr–Sommerfeld quantum conditions for non-separable systems”, *Ann. Phys. (N. Y.)* **4**, 180–188 (1958).
- [98] B. O. Koopman, “Hamiltonian systems and transformations in Hilbert space”, *Proc. Natl. Acad. Sci. USA* **17**, 315 (1931).
- [99] T. Kreilos and B. Eckhardt, “Periodic orbits near onset of chaos in plane Couette flow”, *Chaos* **22**, 047505 (2012).
- [100] Y. Kuramoto and S. Koga, “Anomalous period-doubling bifurcations leading to chemical turbulence”, *Phys. Lett. A* **92**, 1–4 (1982).
- [101] Y. Kuramoto and T. Tsuzuki, “Persistent propagation of concentration waves in dissipative media far from thermal equilibrium”, *Progr. Theor. Phys.* **55**, 365–369 (1976).
- [102] A. Y. Kuznetsova, C. K. A. P. Kuznetsov, and E. Mosekilde, “Catastrophe theoretic classification of nonlinear oscillators”, *Int. J. Bifur. Chaos* **12**, 1241–1266 (2004).
- [103] Y. Lan, *Dynamical Systems Approach to 1 – d Spatiotemporal Chaos – A Cyclist’s View*, PhD thesis (School of Physics, Georgia Inst. of Technology, Atlanta, 2004).
- [104] Y. Lan and P. Cvitanović, “Variational method for finding periodic orbits in a general flow”, *Phys. Rev. E* **69**, 016217 (2004).
- [105] J. G. Leopold and I. Percival, “The semiclassical two-electron atom and the old quantum theory”, *J. Phys. B* **13**, 1037 (1980).
- [106] T.-Y. Li and J. A. Yorke, “Period three implies chaos”, *Amer. Math. Monthly* **82**, 985–992 (1975).
- [107] V. López, P. Boyland, M. T. Heath, and R. D. Moser, “Relative periodic solutions of the Complex Ginzburg-Landau equation”, *SIAM J. Appl. Dyn. Syst.* **4**, 1042 (2006).
- [108] E. N. Lorenz, “Deterministic nonperiodic flow”, *J. Atmos. Sci.* **20**, 130–141 (1963).
- [109] E. N. Lorenz, “Noisy periodicity and reverse bifurcation”, *Annals of the NY Acad. Sci.* **357**, 282–291 (1980).
- [110] A. Lyapunov, “Problème général de la stabilité du mouvement”, *Ann. Math. Studies* **17**, Russian original Kharkow, 1892, 531–534 (1977).
- [111] J. Maurer and A. Libchaber, “Effect of the Prandtl number on the onset of turbulence in liquid 4He”, *J. Physique Lett.* **41**, 515–518 (1980).
- [112] R. M. May, “Simple mathematical models with very complicated dynamics”, *Nature* **261**, 459–467 (1976).
- [113] H. P. McKean, “Selberg’s trace formula as applied to a compact Riemann surface”, *Commun. Pure Appl. Math.* **25**, 225–246 (1972).
- [114] N. Metropolis, M. L. Stein, and P. R. Stein, “On finite limit sets for transformations on the unit interval”, *J. Combin. Theory* **15**, 25–44 (1973).

- [115] J. Milnor and W. Thurston, “Iterated maps of the interval”, in *Dynamical Systems (Maryland 1986-87)*, edited by A. Dold and B. Eckmann (Springer, New York, 1988), pp. 465–563.
- [116] F. C. Moon, *Chaotic Vibrations: An Introduction for Applied Scientists and Engineers* (Wiley, New York, 1987).
- [117] D. W. Moore and E. A. Spiegel, “A thermally excited nonlinear oscillator”, *Astrophys. J.* **143**, 871–887 (1966).
- [118] J. von Neumann, “Zusätze zur Arbeit “Zur Operatorenmethode in der klassischen Mechanik”. (German) [Additions to the work “On operator methods in classical mechanics”]”, *Ann. Math.* **33**, 789–791 (1932).
- [119] J. von Neumann and E. P. Wigner, “Über merkwürdige diskrete Eigenwerte. Über das Verhalten von Eigenwerten bei adiabatischen Prozessen”, *Phys. Zeit.* **30**, 467–470 (1929).
- [120] S. Ostlund, D. Rand, J. Sethna, and E. Siggia, “Universal properties of the transition from quasi-periodicity to chaos in dissipative systems”, *Physica D* **8**, 303–342 (1983).
- [121] E. Ott, “Strange attractors and chaotic motions of dynamical systems”, *Rev. Mod. Phys.* **53**, 655–671 (1981).
- [122] E. Ott, *Chaos and Dynamical Systems* (Cambridge Univ. Press, Cambridge, 2002).
- [123] A. Pais, *Inward Bound: of Matter and Forces in the Physical World* (Oxford Univ. Press, Oxford, 1986).
- [124] A. Pais, *Niels Bohr’s Times, in Physics, Philosophy and Polity* (Oxford Univ. Press, Oxford, 1991).
- [125] W. Parry and M. Pollicott, “An analogue of the prime number theorem for closed orbits of Axiom A flows”, *Ann. Math.* **118**, 573–591 (1983).
- [126] H. Poincaré, *Les Méthodes Nouvelles de la Mécanique Céleste* (Guthier-Villars, Paris, 1899).
- [127] M. Pollicott, “On the rate of mixing of Axiom A flows”, *Inv. Math.* **81**, 413–426 (1985).
- [128] M. Pollicott, “A note on the Artuso-Aurell-Cvitanović approach to the Feigenbaum tangent operator”, *J. Stat. Phys.* **62**, 257–267 (1991).
- [129] D. Rand, S. Ostlund, J. Sethna, and E. D. Siggia, “Universal transition from quasiperiodicity to chaos in dissipative systems”, *Phys. Rev. Lett.* **49**, 132–135 (1982).
- [130] E. Rosenqvist, *Periodic Orbit Theory Beyond Semiclassics: Convergence, Diffraction and \hbar Corrections*, PhD thesis (Copenhagen Univ., Copenhagen, 1995).
- [131] D. Ruelle, “Generalized zeta-functions for Axiom A basic sets”, *Bull. Amer. Math. Soc.* **82**, 153–156 (1976).
- [132] D. Ruelle, “Zeta-functions for expanding maps and Anosov flows”, *Inv. Math.* **34**, 231–242 (1976).

- [133] D. Ruelle, “Locating resonances for Axiom A dynamical systems”, *J. Stat. Phys.* **44**, 281–292 (1986).
- [134] D. Ruelle, “One-dimensional Gibbs states and Axiom A diffeomorphisms”, *J. Diff. Geom.* **25**, 117–137 (1987).
- [135] D. Ruelle, “Resonances for Axiom A flows”, *J. Diff. Geom.* **25**, 99–116 (1987).
- [136] D. Ruelle, “The Deterministic chaos: the science and the fiction”, *Proc. R. Soc. Lond. A* **427**, 241–248 (1990).
- [137] D. Ruelle, *Thermodynamic Formalism: The Mathematical Structure of Equilibrium Statistical Mechanics*, 2nd ed. (Cambridge Univ. Press, Cambridge, 2004).
- [138] H. H. Rugh, “The correlation spectrum for hyperbolic analytic maps”, *Nonlinearity* **5**, 1237 (1992).
- [139] E. Sander and J. A. Yorke, “A period-doubling cascade precedes chaos for planar maps”, *Chaos* **23**, 033113 (2013).
- [140] M. du Sautoy, *Finding Moonshine: A Mathematician’s Journey Through Symmetry* (Harper Collins, 2012).
- [141] I. Segal, “Book review: Alain Connes, Noncommutative geometry”, *Bull. Amer. Math. Soc.* **33**, 459–465 (1996).
- [142] A. Selberg, “Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series”, *J. Indian Math. Soc. (N.S.)* **20**, 47–87 (1956).
- [143] S. J. Shenker, “Scaling behavior in a map of a circle onto itself: Empirical results”, *Physica D* **5**, 405–411 (1982).
- [144] Y. G. Sinai, “Gibbs measures in ergodic theory”, *Russ. Math. Surv.* **27**, 21–69 (1972).
- [145] G. I. Sivashinsky, “Nonlinear analysis of hydrodynamical instability in laminar flames - I. Derivation of basic equations”, *Acta Astronaut.* **4**, 1177–1206 (1977).
- [146] S. Smale, “Differentiable dynamical systems”, *Bull. Amer. Math. Soc.* **73**, 747–817 (1967).
- [147] E. A. Spiegel, “Chaos: A mixed metaphor for turbulence”, *Proc. R. Soc. Lond. A* **A413**, 87 (1987).
- [148] D. Stone, “Einstein’s unknown insight and the problem of quantizing chaos”, *Phys. Today* **58**, 37–43 (2005).
- [149] R. Vilela Mendes, “Critical-point dependence of universality in maps of the interval”, *Phys. Lett. A* **84**, 1–3 (1981).
- [150] E. B. Vul and K. M. Khanin, “The unstable separatrix of Feigenbaum’s fixed-point”, *Russ. Math. Surv.* **37**, 200–201 (1982).
- [151] E. B. Vul, Y. G. Sinai, and K. M. Khanin, “Feigenbaum universality and the thermodynamic formalism”, *Russ. Math. Surv.* **39**, 1–40S (1984).
- [152] H. Wedin and R. R. Kerswell, “Exact coherent structures in pipe flow”, *J. Fluid Mech.* **508**, 333–371 (2004).

- [153] E. W. Weisstein, [Feigenbaum constant](#), 2012.
- [154] E. P. Wigner, *Group Theory and Its Application to the Quantum Mechanics of Atomic Spectra* (Academic, New York, 1931).
- [155] D. Wintgen, K. Richter, and G. Tanner, “The semiclassical helium atom”, [Chaos 2](#), 19–33 (1992).
- [156] S. M. Zoldi and H. S. Greenside, “Spatially localized unstable periodic orbits of a high-dimensional chaotic system”, [Phys. Rev. E 57](#), R2511–R2514 (1998).